SWISS DRUG TESTING

Swiss Drug Testing GmbH Technoparkstrasse 2 8406 Winterthur Switzerland contact@swissdrugtesting.ch www.swissdrugtesting.ch



Medicanna Ltd - EthicaCBD Unit 7 Bude Business Centre EX23 8QN Bude UK

Customer ID: 2000080

Winterthur, 02/13/2021

P10A01

Test Report No.: P10A01

Customer: Medicanna Ltd - EthicaCBD Med - Isolate - 2 Customer sample ID:

Matrix: Oil Lab sample ID:

02/11/2021 Received:

02/11/2021 - 02/13/2021 Analysis period:

Results

Cannabinoid concentrations in mg/kg

Δ8-THC (Δ8-Tetrahydrocannabinol)

Δ 9-THC (Δ 9-Tetrahydrocannabinol)	n.d.	CBG (Cannabigerol)	n.d.
THCA (Tetrahydrocannabinolic acid)	n.d.	CBGA (Cannabigerolic acid)	n.d.
T TIIO		T + 1 0D0	

Total ∆9-THC Total CBG

CBD (Cannabidiol)	105000	CBN (Cannabinol)	n.d.
CBDA (Cannabidiolic acid)	n.d.	CBNA (Cannabinolic acid)	n.d.

Total CBD 105000 Total CBN

CBDV (Cannabidivarin) 133 THCV (Tetrahydrocannabivarin) n.d. CBDVA (Cannabidivarinic acid) n.d. THCVA (Tetrahydrocannabivarinic n.d. acid)

n.d. = not detected, below the limit of detection of 5 mg/kg; < LOQ = below the limit of quantitation of 20 mg/kg for details on sampling and analysis see below

> Dr. Michael Pötzsch CEO, Pharmacist

Analytical details

Sample description

The container was received in good condition already labeled with a number assigned by the customer. The sample was received and stored at ambient temperature. A unique Swiss Drug Testing identification number was assigned and attached to the sample, which was used for further laboratory work and analyses.

Sampling and Analysis of Cannabinoids

A portion of the sample was weighted, extracted into solvent, diluted and spiked with internal deuterated standards. Analysis was conducted using high performance liquid chromatography coupled to tandem mass spectrometry (UHPLC-MS/MS). Quantitation was performed using an external solvent calibration model with certified reference material for all compounds. The method was not validated for the matrices. Matrix effects might have effected the results. Accuracy of the method was assessed by internal quality control. Ratios of two mass transitions and chromatographic retention time assured identity of the compounds. The "total" concentrations of cannabinoid were calculated by applying the following factors: 0.877 for THCA; 0.877 for CBDA, 0.878 for CBGA and 0.876 for CBNA followed by summation with the decarboxylated form.

No part of the text or graphics may be reproduced without the prior written permission of the Swiss Drug Testing GmbH. This document, and all the information, images and intellectual property rights in it belong to the Swiss Drug Testing GmbH. With sending the sample to our lab, the customer accepted our terms of services, which are published on www.swissdrugtesting.ch.